arXiv:1202.5351v1 [math.PR] 24 Feb 2012

BOOTSTRAP PERCOLATION ON THE HAMMING TORUS

CHRISTOPHER HOFFMAN AND JAMES PFEIFFER

Abstract. We consider bootstrap percolation on the Hamming torus and calculate the threshold functions for subgraphs of different dimensions to become open. We show that the threshold function for 1-dimensional subgraphs is distinct from higher dimensional subgraphs, while the threshold functions for i dimensional subgraphs, $i \geq 2$, are very closely bunched.

1. Introduction

Bootstrap percolation is a model for various physical phenomena including crack formation, clustering, sandpiles, and the Ising model of ferromagnetism which was introduced in 1979 by Chalupa, Leith and Reich [7]. For more applications and background see J. Adler and U. Lev, Bootstrap Percolation: visualizations and applications [1].

Given a graph $G = (V, E)$, bootstrap percolation with threshold T is defined as follows. Given an initial configuration $\omega \in \{0,1\}^V$, bootstrap percolation is an increasing sequence $\omega = \omega_0, \omega_1, \ldots$, defined by

$$\omega_{i+1}(v) = \begin{cases} 1 & \text{if } \omega_i(v) = 1 \text{ or } \sum_{w \sim v} \omega_i(w) \geq T \\ 0 & \text{else.} \end{cases}$$

The initial configuration ω is random; $\{\omega(v) : v \in V\}$ is a collection of i.i.d. Bernoulli random variables with parameter $p = p(n)$. The central question in bootstrap percolation is to determine the critical probability $p_c = p_c(n)$ such that

$$\mathbb{P}_{p_c}(\omega_\infty = 1) = 1/2.$$

Bootstrap percolation has been studied on various graphs. The first results were by van Enter and Schonmann, who proved that the critical probability for $\mathbb{Z}^d$ is either 0 or 1 according to whether $T \leq d$ or $T > d$ [10]. Aizenman and Lebowitz considered bootstrap percolation on the finite grid $[n]^d$ and $T = 2$, and showed that the critical probability is on the order of $(\frac{1}{\log n})^{d-1}$ [2]. Later, Cerf and Cirillo for $d = T = 3$, and Cerf and Manzo for general d and T showed that the critical probability is on the order of $(\log_{T-1} n)^{-d+T-1}$, with $\log_{T-1}$ denoting the iterated logarithm [6] [5].

The first sharp description of the critical probability was due to Holroyd, who showed that for $d = T = 2$, $p = \frac{\pi^2}{18 \log n} + o\left(\frac{1}{\log n}\right)$ [8].This result is highly celebrated as it revealed different results than simulations indicated due to a "crossover" phenomenon. Later Balogh, Bollobas, and Morris determined the sharp description of the critical probability for $d = T = 3$, and Balogh, Bollobas, Duminil-Copin, and Morris found the critical probability for all d and T [3] [4].

Besides varying the dimension of the lattice or its threshold, one can also vary what the neighborhood of a point is. In [6] Holroyd, Liggett and Romik study k-boostrap percolation in two dimensions, in which the neighborhood of a point consists of the $k - 1$ points in each axis-parallel direction, with the percolation threshold $T = k$. In this case they determine that the critical probability is asymptotically $\frac{\pi^2}{3k(k+1)\log n}$ [9].

In this paper we consider bootstrap percolation on the Hamming torus with vertex set $V = [n]^d$, where v and w are adjacent iff $v - w$ has exactly one nonzero coordinate. This model is similar in spirit to taking $k = \infty$ of the above model. That is, the neighborhood of a point v is the union of the axis-parallel lines through v (although the threshold T remains fixed). With such large neighborhoods, we get a qualitatively different critical probability than in the above results. Instead of having $p_c = O(\frac{1}{\log(n)})$ as we saw when $d = T = 2$, it is easy to see that the critical probabilities in our model are of the order $n^{-1-\beta}$ for $\beta > 0$. In addition, the models above exhibit a single sharp threshold, below which growth stopped quickly and was limited to small subgraphs; and above which the entire graph became open with high probability. When

$d = 2$ we see a similar phenomenon. We find a critical probability p_c such that when $p = o(p_c)$, then with high probability $\omega_0 = \omega_\infty \ll 1$; and when p grows faster than p_c, then with high probability $\omega_\infty = 1$. We also determine the exact function p_c in this case.

However, in dimensions 3 and greater we find two distinct thresholds. Below the first threshold, growth is small. Between the two thresholds, there is a high probability of lines (one-dimensional subgraphs) becoming open, but asymptotically almost surely there are no two dimensional subgraphs that are eventually open and the graph itself does not percolate. Above the second threshold, 2-dimensional subgraphs become open with high probability. We can also define a similar such threshold for subspaces of dimension i, $1 \leq i \leq d$, It remains an open question to determine if the thresholds for dimension i are distinct for $i \geq 2$. In the next section we give rigorous (but not the strongest) statement of our results. In the subsequent sections we state (and prove) our results in their full generality.

2. Statement of Results

Let F be a family of subsets of $[n]^d$. Then

$$\mathbb{P}_p(\exists f \in F:\ \omega_\infty|_f = 1)$$

is an increasing function in p. For F_i the collection of i-dimensional subgraphs of G, there exists a threshold function $p_c(i, d)$ such that

$$\mathbb{P}_{p_c(i,d)}(\exists f \in F_i:\ \omega_\infty|_f = 1) = .5.$$

If $\omega_i(v) = 1$ we say v is open at step i, and a set $S \subseteq V$ is open if each $v \in S$ is open.

The smaller that i, is the easier it is to calculate $p_c(i, d)$. For $i = 0$ we have a second critical probability. We would like to define $p_c^*(0, d)$ to be the threshold function for $\omega_\infty \neq \omega_0$. Unfortunately, this is not an increasing event. So instead we define $p_c^*(0, d)$ to be the threshold function for the event

$$\text{Above Threshold} = \left\{ \exists v : \sum_{w \sim v} \omega_w \geq T \right\}.$$

We conjecture that for all $i \geq 1$ and d, p_c has the following form.

Conjecture 2.1. *For all i and d and n sufficiently large*

$$p_c(i, d) = n^{-1-\frac{c_1(i,d)}{T}-\frac{c_{3/2}(i,d)}{T^{3/2}}+o(T^{-3/2})}.$$

We make substantial progress to proving that this is the case for all i and d, but in general we are only able to prove that if n is sufficinetly large with T then

$$p_c(i, d) = n^{-1-\frac{c_1(i,d)}{T}-\theta(T^{-3/2})}.$$

In general for fixed n, d and i we get bounds on the critical p, but the precise bounds that we get are quite messy to state. In the rest of this section we put all of our results in a common form. Many of the theorems have are given in a stronger form in the following sections.

When $d = 2$ we get our most precise answer.

Theorem 2.1. *Let $d = 2$, $k = \lceil T/2 \rceil > 1$ and $p = an^{-1-\frac{1}{k}}$. Then*

$$\mathbb{P}(\omega_\infty = 1) \to \begin{cases} 1 - e^{-2a^k/k!} & \textit{if } T \textit{ is odd,} \\ (1 - e^{-a^k/k!})^2 & \textit{if } T \textit{ is even.} \end{cases}$$

Thus

$$p_c(2, 2) = p_c(1, 2) = p_c^*(0, 2) = n^{-1-\frac{2}{T}+o(T^{-3/2})},$$

Furthermore

$$\mathbb{P}\big(\{\omega_\infty \neq \omega_0\} \setminus \{\omega_\infty = 1\}\big) = o(1).$$

We are also able to calculate $p_c^*(0, d)$ and $p_c(1, d)$ for all $d \geq 2$ quite precisely as well.

Theorem 2.2. *Let* $p = f(n)n^{-1-\frac{d}{T}}$ *and* $d, T \geq 3$. *If* $f(n) \to 0$ *then*

$$\mathbb{P}(\textit{Above Threshold}) \to 0$$

and if $f(n) \to \infty$ *then*

$$\mathbb{P}(\exists \textit{ a line } L \textit{ such that } \omega_\infty|_L = 1) \to 1.$$

Thus $c_1(1, d) = d$ *and* $c_{3/2}(1, d) = 0$.

We can also get good bounds on $p_c(2, d)$, the threshold for existence of two dimensional subspaces in the final configuration.

Theorem 2.3. *Fix* d *and* T. *For* n *sufficiently large*

$$n^{-1-\frac{2}{T}-\frac{4d^2+2}{T^{3/2}}} \leq p_c(2, d) \leq n^{-1-\frac{2}{T}-\frac{1}{T^{3/2}}}.$$

There has been no attempt to optimize the constants 1 and $4d^2 + 2$ in the above theorem.

The higher the dimensions i and d become the more difficult it becomes to calculate $p_c(i, d)$. However Theorems 2.2 and 2.3 are sufficient for us to get bounds on $p_c(i, d)$ for all $i, d \geq 2$.

Theorem 2.4. *For all* i *and* d *and sufficiently large* n

$$p_c(i, d) = n^{-1-\frac{c_1(i,d)}{T}-\theta(T^{-3/2})}.$$

Proof. It is easy to see that $p_c(i, d)$ is nondecreasing in i and decreasing in d. Also $p_c(d, d)$ is decreasing in d. To see this last inequality note that when $n \geq 3T$ and $d = j + 1$

$$\mathbb{P}_{p_c(j,j)}(\exists \text{ at least } T\ i \text{ such that } \omega_\infty|_{(i,*,*,\cdots)} = 1) > 1/2.$$

The event on the left hand side implies that $\omega_\infty = 1$ and thus

$$p_c(j + 1, j + 1) \leq p_c(j, j)$$

and inductively

$$p_c(d, d) \leq p_c(3, 3).$$

So

$$p_c(2, d) \leq p_c(i, d) \leq p_c(d, d) \leq p_c(3, 3).$$

By Theorem 2.3

$$p_c(2, 3) \leq n^{-1-\frac{2}{T}-\frac{1+o(1)}{T^{3/2}}}.$$

By coupling it is easy to see that ω chosen when $p = 10Tp_c(2, 3)$ stochastically dominates the union of $10T$ independent ω' chosen with $p = p_c(2, 3)$. Then by the definition of $p_c(2, 3)$

$$\mathbb{P}_{10Tp_c(2,3)}(\exists \text{ at least } T\ i \text{ such that } \omega_\infty|_{(i,*,*)} = 1) > 1/2.$$

The event on the left hand side implies $\omega|_\infty = 1$ and thus

$$p_c(3, 3) \leq 10Tp_c(2, 3).$$

And putting this all together for all $d \geq 3$ and $2 \leq i \leq d$

$$n^{-1-\frac{2}{T}-\frac{40+o(1)}{T^{3/2}}} \leq p_c(2, d) \leq p_c(i, d) \leq p_c(d, d) \leq p_c(3, 3) \leq 10Tp_c(2, 3) \leq n^{-1-\frac{2}{T}-\frac{2+o(1)}{T^{3/2}}}$$

which is the desired result. ☐

Remark 2.5. The above theorems hold for n sufficiently large. We could calculate how large n needs to be. Doing so involves a close look at the constants in the following sections. These constants depend on T. In order to get the all the proofs to work we must choose n to be roughly $e^{T^{3/2}}$. As our calculations were optimized for ease of calculation, not to get the best possible bounds, we do not worry about for exactly which values of n our theorems apply.

The rest of the paper is organized as follows. In Section 3 we find the threshold function for the two-dimensional case. In Section 4 we establish the threshold function for a line to become open. In Section 5 we establish an upper bound on the threshold function for a plane to become open and in Section 6 we establish a lower bound the threshold function for a plane to become open.

3. 2-DIMENSIONAL RESULTS

In the two-dimensional case we can describe the limiting behavior exactly as $n \to \infty$. Let $k = \lceil T/2 \rceil$ and $p = an^{-1-\frac{1}{k}}$ for some constant a. Also assume $k > 1$; the cases $T = 1$ and $T = 2$ are easy to work out separately. (For $T = 1$, $\omega_\infty = 1$ if and only if $\omega_0 \neq 0$; for $T = 2$, $\omega_\infty = 1$ asymptotically if and only if ω_0 has at least two points open).

Lemma 3.1. *Let $k = \lceil T/2 \rceil$ and $p = an^{-1-\frac{1}{k}}$. With probability going to 1, there are no lines with at least $k+1$ points initially open.*

Proof. The probability that a given line has $k+1$ points initially open is bounded above by $\binom{n}{k+1}p^{k+1} \leq a^{k+1}n^{-1-\frac{1}{k}}$. Since there are $2n$ lines the union bound shows that asymptotically almost surely, none of them have $k+1$ points. $\square$

Lemma 3.2. *Let $k = \lceil T/2 \rceil$ and $p = an^{-1-\frac{1}{k}}$. Fix constants A, B. With probability going to 1, there are at least A horizontal (vertical) lines with $k-1$ points initially open. Their $k-1$ open sites may be taken to avoid any B fixed vertical (horizontal) lines.*

Proof. The probability that a given horizontal line satisfies the conditions is bounded below by $\binom{n-B}{k-1}p^{k-1}(1-p)^{n-k+1} = O(n^{-1+\frac{1}{k}})$. There are n horizontal lines so the mean number of such lines goes to ∞. Therefore the probability that there are less than A of them goes to 0. $\square$

Let E_{horiz} be the event that some horizontal line has at least k points initially open, and E_{vert} the corresponding event for vertical lines.

Lemma 3.3. *Let $E_{horiz} \circ E_{vert}$ be the event that E_{horiz} and E_{vert} occur disjointly. Then*

$$P((E_{horiz} \cap E_{vert}) \setminus (E_{horiz} \circ E_{vert})) \to 0.$$

Furthermore,

$$P(E_{horiz} \cap E_{vert}) \to P(E_{horiz})P(E_{vert}) = (1 - e^{-a^k/k!})^2$$

and

$$P(E_{horiz} \cup E_{vert}) \to 1 - ((e^{-a^k/k!})^2.$$

Proof. Since E_{horiz} and E_{vert} are increasing events, by the FKG inequality we have $P(E_{\text{horiz}} \cap E_{\text{vert}}) \geq P(E_{\text{horiz}})P(E_{\text{vert}})$. By the BK inequality we have $P(E_{\text{horiz}})P(E_{\text{vert}}) \geq P(E_{\text{horiz}} \circ E_{\text{vert}})$. But $(E_{\text{horiz}} \cap E_{\text{vert}}) \setminus (E_{\text{horiz}} \circ E_{\text{vert}})$ happens only if some point is open but each crossline has exactly $k-1$ open. The probability of which is bounded (by the union bound) by

$$n^2 p \left(\binom{n}{k-1} p^{k-1} \right)^2 = O(n^{-1+\frac{1}{k}}) \to 0.$$

Thus asymptotically, $P(E_{\text{horiz}} \cap E_{\text{vert}}) \to P(E_{\text{horiz}})P(E_{\text{vert}}) = P(E_{\text{horiz}})^2$. But E_{horiz} is exactly the union of n independent events with probability asymptotically $\frac{a^k}{k!}n^{-1}$; so by Poisson approximation, the chance of its success is asymptotically $1 - e^{-a^k/k!}$.

$\square$

Let F be the event that the entire graph becomes open; i.e. $F = \{\omega_\infty = 1\}$.

Lemma 3.4. *If T is even, $P(F \Delta (E_{horiz} \cap E_{vert})) \to 0$, while if T is odd,*

$$P(F \Delta (E_{horiz} \cup E_{vert})) \to 0.$$

Proof. If T is odd no percolation can take place unless there is some line with at least k points initially open. So $F \subseteq E_{\text{horiz}} \cup E_{\text{vert}}$. If T is even, then by the pigeonhole principle and Lemma 3.1, $P(F \setminus (E_{\text{horiz}} \cup E_{\text{vert}})) \to 0$.

Now suppose T is odd and $E_{\text{horiz}} \cup E_{\text{vert}}$ occurs. Then some line L has k points open initially. By Lemma 3.2 there are T lines $\{J_i\}$ parallel to L, each with $k-1$ points open. Again by Lemma 3.2 there are T lines $\{H_j\}$ crossing L, each with $k-1$ points open, and with their open sites avoiding the J_i and L. For each point of intersection, v, of H_j and L, v has T neighbors open: $k-1$ from H_j and k from L. Thus v goes on. Since L has T points open, L becomes open. Now for each point w of intersection of H_j and J_i, w sees 1 open neighbor on L, and $k-1$ open neighbors each on H_j and J_i, for a total of T. Thus w becomes open. Since there are T such points on each H_i and J_i, these $2T$ lines become open, and the entire graph becomes open.

Finally, suppose T is even and $E_{\text{horiz}} \cap E_{\text{vert}}$ occurs. With probability going to 1, we may assume $E_{\text{horiz}} \circ E_{\text{vert}}$ occurs by Lemma 3.3. That is, there is a horizontal line L and a vertical line L', each with k points initially open, excluding their point of intersection. This point of intersection becomes open. By Lemma 3.2 there are T lines in each direction, J_i and J_i', each with $k-1$ initially open points, and such that no open sites on one line meet any other lines. Then every point of intersection between L and J_i', and between L' and J_i, sees $T = (k+1) + (k-1)$ open sites, so it becomes open. Then L and L' have T open sites, so they become open. Then every point of intersection of a J_i with a J_i' sees $2 + 2(k-1) = T$ open sites, so becomes open. Now as in the odd-T case, the entire graph becomes open.

$\square$

Theorem 3.5. *Let $T \geq 3$, $p = an^{-1-\frac{1}{k}}$ where $k = \lceil \frac{T}{2} \rceil$. As $n \to \infty$,*

$$\lim_{n\to\infty} \mathbb{P}(F) = \begin{cases} (1 - e^{-a^k/k!})^2 & \text{if } T \text{ is even,} \\ 1 - (e^{-a^k/k!})^2 & \text{if } T \text{ is odd.} \end{cases}$$

Proof. This follows from Lemmas 3.3 and 3.4. $\square$

4. One-dimensional subgraphs

One mechanism for a line L to become open is for it to start with $\alpha \leq T$ vertices initially open, and to intersect one line with $T-\alpha$ sites (not on L) initially open and T other lines, each with $T-\alpha-1$ sites (not on L) initially open. Then after one step, L has $\alpha+1$ points open, and after two steps, T points open. After three steps L is completely open. We now introduce some notation to use this method to get an upper bound on the threshold function for lines.

For a set $S \subset V$ and $x \in \mathbf{N}$, let $\text{Initial}\,(S, \geq x)$ be the event that the set S has at least x points initially open, i.e.

$$\text{Initial}\,(S, \geq x) = \left\{ \sum_{v \in S} \omega_0(v) \geq x \right\}.$$

For a point $v \in V$, let $P_{1,2}(v)$ be the e_1, e_2-parallel plane through v:

$$P_{1,2}(v) = \{(a_1, a_2, v_3, v_4, \ldots, v_d) : \ a_1, a_2 \in [n]\}.$$

Let $L_2(v)$ be the e_2-parallel line through v:

$$L_2(v) = \{(v_1, a_2, v_3, v_4, \ldots, v_d) : \ a_2 \in [n]\}.$$

For any e_1-parallel line L, define

$$L_l = \{w \in L, w_1 < n/3\}, \quad L_m = \{w \in L, n/3 \leq w_1 \leq 2n/3\}, \quad \text{and} \quad L_r = \{w \in L, w_1 > 2n/3\}$$

to be the left, middle and right thirds of L. Define

$$\text{Cross Lines}_m(L) = \left\{ \sum_{v \in L_m} \mathbf{1}_{\text{Initial}(L_2(v), \geq T-\alpha)} \geq 1 \right\}$$

$$\text{Cross Lines}_r(L) = \left\{ \sum_{v \in L_r} \mathbf{1}_{\text{Initial}(L_2(v), \geq T-\alpha-1)} \geq T \right\}$$

and

$$F_L = \text{Initial}\,(L_l, \geq \alpha) \bigcap \text{Cross Lines}_m(L) \bigcap \text{Cross Lines}_r(L).$$

Notice that the event F_L depends only on the sites in $P_{1,2}(v)$ for any $v \in L$. Also note that

$$\text{Cross Lines}_m(L) = \text{Cross Lines}_m(L') \quad \text{and} \quad \text{Cross Lines}_r(L) = \text{Cross Lines}_r(L')$$

for any e_1 parallel lines L and L' that lie in a common e_1, e_2 plane. Finally, note that $\text{Initial}\,(L_l, \geq \alpha)$, $\text{Cross Lines}_m(L)$, and $\text{Cross Lines}_r(L)$ are independent, and $\text{Initial}\,(L_l, \geq \alpha)$ and $\text{Initial}\,(L_l', \geq \alpha)$ are independent.



FIGURE 1. An instance of the event F_L. Here $T = 6$, $\alpha = 3$. After one step the intersection of lines L and L' becomes open so L has $\alpha + 1$ vertices open. At step 2 the T intersections with L and the other T vertical lines become open. At step 3 all of L becomes open.

Lemma 4.1. *If L is a line parallel to the e_1 axis and F_L occurs, then the entire line L is open after three steps.*

Proof. If $v \in L_m$ and the event

$$\text{Initial}\left(L_l, \geq \alpha\right) \bigcap \text{Initial}\left(L_2(v), \geq T-\alpha\right)$$

occurs then v has at least T neighbors initially open and v is open after one step. Then there are at least $\alpha + 1$ sites open in $L_l \cup L_m$ after one step. Thus if F_L occurs there are at least T vertices in L_r that are open after two steps. Thus the entire line L is open after three steps. ☐

Remark 4.2. F_L is chosen due to the independence of the events involved, which makes computations easier. A more natural definition would not restrict the orientations of the lines, or the left/middle/right sections thereof. By restricting the event in this way we reduce the probability of F_L by a constant factor, independent of n.

We set $\alpha = \left\lceil \frac{(d-1)T}{d} \right\rceil - 1$ and $p = n^{-1-\frac{d}{T}} f(n)$, where $f(n)$ is any function such that $f(n) \to \infty$. We will show that in this regime some line becomes open asymptotically almost surely.

Lemma 4.3. *Fix $v \in V$ and $T, d \geq 3$. Let $p = n^{-1-\frac{d}{T}} f(n)$ where $f(n) \to \infty$. Then for any $c > 0$, the probability that there exists an e_1-parallel line L in $P_{1,2}(v)$ such that F_L occurs is at least cn^{2-d} for n sufficiently large.*

Proof. First note that when $T \geq d \geq 3$ then $\alpha d/T \geq 1$ as

$$\alpha d/T \geq \left(\frac{(d-1)T}{d} - 1\right) \frac{d}{T} \geq d - 1 - d/T.$$

The right hand side is greater than 1 except if $d = T = 3$. For now we assume that at least one of d and T are greater than 3.

The three events that define F_L depend on disjoint sets of sites, so they are independent. Also only the first depends on L. For any L

$$\begin{aligned}\mathbb{P}(\text{Initial}\,(L_l,\geq\alpha)) &\geq \binom{n/3}{\alpha}p^{\alpha}(1-p)^{n/3-\alpha}\\ &\geq c_1(np)^{\alpha}\\ &\geq c_1(f(n)n^{-d/T})^{\alpha}.\end{aligned}$$

Provided that $f(n)$ does not grow too quickly the right hand side is much smaller than $1/n$ so we get

$$\mathbb{P}\big(\exists L \text{ such that Initial}\,(L_l,\geq\alpha) \text{ occurs }\big)\geq .5c_1n(f(n)n^{-d/T})^{\alpha}.$$

The second probability is bounded below by the probability that a

$$\text{Binomial}\left(n/3,\binom{n}{T-\alpha}p^{T-\alpha}(1-p)^{n-(T-\alpha)}\right)$$

random variable is at least one. If $f(n)$ doesn't grow too fast then $(n/3)p^{T-\alpha}(1-p)^{n-(T-\alpha)}<2$ and

$$\begin{aligned}\mathbb{P}(\text{Cross Lines}_m(L)) &\geq (1/10)(n/3)\binom{n}{T-\alpha}p^{T-\alpha}(1-p)^{n-(T-\alpha)}\\ &\geq c_2n(np)^{T-\alpha}\\ &\geq c_2n(f(n)n^{-d/T})^{T-\alpha}.\end{aligned}$$

The third probability is bounded below by the probability that a

$$\text{Binomial}\left(n/3,\binom{n}{T-\alpha-1}p^{T-\alpha-1}(1-p)^{n-(T-\alpha-1)}\right)$$

random variable is at least T. Its expected value is

$$\begin{aligned}(n/3)\binom{n}{T-\alpha-1}p^{T-\alpha-1}(1-p)^{n-(T-\alpha-1)} &\geq c_3n(np)^{T-\alpha-1}\\ &\geq c_3f(n)^{T-\alpha-1}n^{1-d+(\alpha+1)\frac{d}{T}}\\ &\geq c_3f(n)^{T-\alpha-1}.\end{aligned}$$

As this goes to infinity the probability that it is bigger than any fixed number goes to 1.
Multiplying together the probabilities we get for any c

$$\begin{aligned}&\mathbb{P}(\exists\ L \text{ in } P_{1,2}(v) \text{ such that that } F_L \text{ occurs})\\ &\quad= \mathbb{P}(\exists L \text{ such that Initial}\,(L_l,\geq\alpha))\mathbb{P}(\text{Cross Lines}_m(L))\mathbb{P}(\text{Cross Lines}_r(L))\\ &\quad\geq .5c_1n(f(n)n^{-d/T})^{\alpha}c_2n(f(n)n^{-d/T})^{T-\alpha}(1/2)\\ &\quad\geq c_3n^2(f(n)n^{-d/T})^{T}\\ &\quad\geq cn^{2-d}.\end{aligned}$$

As the probability of this event is monotone in $f(n)$ we can remove our restriction that $f(n)$ does not grow too fast, and the lemma is true. The case when $d=T=3$ is almost identical except the roles of the first and second events are reversed. This happens because $\alpha=1<T-\alpha=2$. This is the only choice of d and T for which $\alpha<T-\alpha$. We leave the details to the reader. □

Theorem 4.4. *Suppose* $p=n^{-1-\frac{d}{T}}f(n)$ *with* $f(n)\to\infty$. *Then* $\mathbb{P}(\cup_L F_L)\to 1$ *as* $n\to\infty$, *where the union is taken over all* e_1 *parallel lines. Thus with probability going to 1, some line becomes open after three steps.*

Proof. We can choose n^{d-2} distinct v_i such that $P_{1,2}(v_i)$ are disjoint. Then the events that there exist L in $P_{1,2}(v_i)$ where F_L occurs are independent. Thus $\mathbb{P}(\cup_L F_L)$ is bounded below by the probability that a

$$\text{Binomial}(n^{d-2}, \mathbb{P}(\exists L \text{ in } P_{1,2}(v_i) \text{ such that } F_L \text{ occurs}))$$

is at least one. The expected value of the latter is

$$n^{d-2}\mathbb{P}(\exists L \text{ in } P_{1,2}(v_i) \text{ such that } F_L \text{ occurs}) \geq n^{d-2}cn^{2-d} = c$$

for any fixed c. Thus $\mathbb{P}(\cup_L F_L) \to 1$. ☐

Theorem 4.5. *Let $p = n^{-1-d/T}f(n)$. If $f(n) \to 0$, then $\mathbb{P}$(Above Threshold) $\to 0$.*

Proof. We compute the expected number of points that become open at the first stage. First let $f(n) \to 0$. Using the union bound,

$$\begin{aligned} \mathbb{P}(\text{Above Threshold}) &\leq \sum_{v \in V} \mathbb{P}\left(\sum_{w \sim v} \omega_0(w) \geq T\right) \\ &= n^d \mathbb{P}\left(\sum_{w \sim v} \omega_0(w) \geq T\right) \\ &\leq n^d \binom{n}{T} p^T \\ &\leq f(n)^T \end{aligned}$$

which approaches 0 since $f(n) \to 0$ as $n \to \infty$. ☐

Proof of Thm 2.2. Combining Theorems 4.4 and 4.5 proves the result. The assuption that $T \geq 3$ is needed so that $\alpha \geq 1$. ☐

5. Sufficiency for planes

As with lines, we will give a critical probability and an event F_P such that if F_P occurs, then the plane P becomes open, and such that $\cup_P F_P$ occurs asymptotically almost surely. Again we will restrict the event F_P for ease of calculation.

Fix a point $v = (v_1, v_2, \ldots, v_n)$. As before, define $P = P_{1,2}(v)$ to be the e_1, e_2 plane containing v. Split P into four quadrants Q_i, $i = 1, \ldots, 4$, with the same usual labeling as the quadrants in $\mathbb{R}^2$ (with the origin $(n/2, n/2)$.) Also, for $b, c \in [n]$, define lines $L_1(b)$

$$L_1(b) = \{(e, b, v_3, \ldots, v_d) : \ e \in [n]\},$$

and quarter lines $QL_2(c)$

$$QL_2(c) = \{(c, e, v_3, \ldots, v_d) : 3n/4 < e \leq n\}.$$

Fix some $\alpha, 1 \leq \alpha \leq T$ - we will determine α later. Let A_P be the event that in the bottom half of quadrant 1, at least α e_1-parallel lines have $\frac{T+\alpha}{2}$ points initially open:

$$A_P = \left\{ \sum_{n/2 < b \leq 3n/4} 1_{\text{Initial}\left(L_1(b) \cap Q_1, \geq \frac{T+\alpha}{2}\right)} \geq \alpha \right\},$$

and B_P be the event that in quadrant 2 there are at least T e_2 parallel quarter lines with $\frac{T-\alpha}{2}$ vertices initially open and in quadrants 3 and 4 there are at least T e_1-parallel lines that have $\frac{T-\alpha}{2}$ vertices initially open:

$$B_P = \left\{ \sum_{c < n/2} 1_{\text{Initial}\left(QL_2(c), \geq \frac{T-\alpha}{2}\right)} \geq T \right\} \bigcap \left\{ \sum_{b \leq n/2} 1_{\text{Initial}\left(L_1(b), \geq \frac{T-\alpha}{2}\right)} \geq T \right\}.$$

Let $F_P = A_P \cap B_P$.

Lemma 5.1. *If P is a plane parallel to the e_1e_2 plane and F_P occurs, then the entire plane P is open after four steps.*



FIGURE 2. An instance of the event F_P. Here $T = 6$, $\alpha = 2$. After one step the T half lines in Q_2 all have $(T+\alpha)/2$ vertices open. After 2 steps the T lines in $Q_3 \cup Q_4$ all have at least T vertices open. After 3 steps those T lines are all open and after 4 steps the entire plane P is open.

Proof. After one step, each intersection of a $Q_1 - Q_2$ pair sees $\frac{T+\alpha}{2} + \frac{T-\alpha}{2} = T$ open points, so such vertices become open. Then in step 2 a $(Q_3 \cup Q_4) - Q_2$ intersection is either open or has $\frac{T+\alpha}{2} + \frac{T-\alpha}{2} = T$ neighbors open so it becomes open. There are T of these intersections on each $Q_3 \cup Q_4$ line, so at step 3 we see T lines become open, so the entire plane P becomes open by step 4. □

We set $p = n^{-1-\frac{2}{T-\alpha}}$. We will determine α later, such that the following holds:

$$-2\alpha^2 - (d-2)\alpha + (d-2)T \geq 0. \tag{5.1}$$

First we work on showing that the probability of B_P is close to one.

Lemma 5.2. *Let α be as above and $p = n^{-1-\frac{2}{T-\alpha}} f(n)$ with $f(n) \to \infty$. Then for a given quarter line QL,*

$$n \cdot \mathbb{P}\left(\textit{Initial}\left(QL, \geq \frac{T-\alpha}{2}\right)\right) \to \infty.$$

Proof. As long as $f(n)$ does not grow too quickly, we have for some $c > 0$ that

$$\begin{aligned}\mathbb{P}\left(\text{Initial}\left(QL, \geq \frac{T-\alpha}{2}\right)\right) &\geq \binom{n/4}{\frac{T-\alpha}{2}} p^{\frac{T-\alpha}{2}} (1-p)^{n-\frac{T-\alpha}{2}} \\ &\geq c(np)^{\frac{T-\alpha}{2}} \\ &= cn^{-1} f(n)^{\frac{T-\alpha}{2}}\end{aligned}$$

Multiplying by n completes the proof. □

Remark 5.3. In the proof of the preceding lemma, as well as several following, we assume that $f(n)$ does not grow too quickly. This means that terms of the form $(1-p)^n$ converge to 1, or at least to positive constants. But since the results of the lemma are clearly monotone in $f(n)$, we can then remove the restriction that $f(n)$ does not grow too quickly.

Lemma 5.4. *Let $p = n^{-1-\frac{2}{T-\alpha}} f(n)$ and suppose $f(n) \to \infty$. Then for any given plane P, $\mathbb{P}(B_P) \to 1$.*

Proof. B_P is the intersection of two independent events (referencing quadrant 2 and quadrants 3,4). We will show that the Q_2 event occurs with probability going to 1. This clearly implies that the probability of the $Q_3 \cup Q_4$ event, and therefore of B_P, goes to 1 as well. Since

$$X = \sum_{QL_2 \subset Q_2} \mathbf{1}_{\text{Initial}(QL_2, \geq \frac{T-\alpha}{2})}$$

is a binomial random variable with $n/2$ trials and $\geq cn^{-1}f(n)^{\frac{T-\alpha}{2}}$ probability of success, its means approach ∞. Thus we may use Poisson approximation to conclude that the chance that $X < T$ converges to 0. $\square$

Now we calculate the probability of A_P.

Lemma 5.5. *Let $p = n^{-1-\frac{2}{T-\alpha}}f(n)$ and suppose $f(n) \to \infty$. Then for a given (length-$n/2$) line L,*

$$n^{\frac{T+\alpha}{T-\alpha}} \cdot \mathbb{P}\left(\mathit{Initial}\left(L, \geq \frac{T+\alpha}{2}\right)\right) \to \infty.$$

Proof. As long as $f(n)$ does not grow too quickly, we have for some $c > 0$ that

$$\begin{aligned}\mathbb{P}\left(\text{Initial}\left(L, \geq \frac{T+\alpha}{2}\right)\right) &\geq \binom{n/2}{\frac{T+\alpha}{2}} p^{\frac{T+\alpha}{2}}(1-p)^{n-\frac{T+\alpha}{2}} \\ &\geq c(np)^{\frac{T+\alpha}{2}} \\ &= cn^{-\frac{T+\alpha}{T-\alpha}}f(n)^{\frac{T+\alpha}{2}}\end{aligned}$$

Multiplying by $n^{\frac{T+\alpha}{T-\alpha}}$ completes the proof. $\square$

Now we are finally ready to choose α. Let it be any integer satisfying

$$-2\alpha^2 - (d-2)\alpha + (d-2)T \geq 0 \tag{5.2}$$

Lemma 5.6. *Let $p = n^{-1-\frac{2}{T-\alpha}}f(n)$ where $f(n) \to \infty$. If (5.2) is satisfied then $\mathbb{P}(\bigcup_P A_P) \to 1$.*

Proof. Fix a plane P. Since

$$\sum_{n/2 < b < 3n/4} \mathbf{1}_{\text{Initial}(L_1(b) \cap Q_1, \geq \frac{T+\alpha}{2})}$$

is a binomial random variable with $n/4$ trials, by Lemma 5.5 we have

$$\begin{aligned}\mathbb{P}(A_P) &\geq \binom{n/2}{\alpha}\left(cf(n)^{\frac{T+\alpha}{2}}n^{-\frac{T+\alpha}{T-\alpha}}\right)^{\alpha}\left(1 - cf(n)^{\frac{T+\alpha}{2}}n^{-\frac{T+\alpha}{T-\alpha}}\right)^{n-\alpha} \\ &\geq c'f(n)^{\frac{T+\alpha}{2}\alpha}n^{\alpha}n^{-\frac{T+\alpha}{T-\alpha}\alpha}\end{aligned}$$

Therefore, $\sum_P \mathbf{1}_{A_P}$ is a binomial random variable with means

$$n^{d-2}\mathbb{P}(A_P) \geq c'f(n)^{\frac{T+\alpha}{2}\alpha}n^{\alpha}n^{-\frac{T+\alpha}{T-\alpha}\alpha}n^{d-2}.$$

The exponent of n is nonnegative because of (5.2) Since $f(n) \to \infty$, we can apply Poisson approximation to conclude

$$\mathbb{P}\left(\bigcup_P A_P\right) \to 1.$$

$\square$

Theorem 5.7. *Let $p = n^{-1-\frac{2}{T-\alpha}}f(n)$ and suppose $f(n) \to \infty$. Then $\mathbb{P}\left(\bigcup_P F_P\right) \to 1$.*

Proof. Consider a family of n^{d-2} disjoint planes. Then the events $\{A_P\}$ are independent of the events $\{B_P\}$ as they depend on disjoint sets of vertices.

$$\begin{aligned}\mathbb{P}\left(\bigcup_P F_P\right) &= \mathbb{P}\left(\bigcup_P F_P \;\middle|\; \bigcup_P A_P\right)\mathbb{P}\left(\bigcup_P A_P\right)\\ &\geq \mathbb{P}(B_P)\mathbb{P}\left(\bigcup_P A_P\right)\\ &\to 1.\end{aligned}$$

□

6. Necessity for planes

In the previous sections we have seen several ways for a vertex to become open. Among these are

(1) v is initially open.
(2) The neighborhood of v has at most T vertices initially open. In this case v becomes open by time 1. And finally
(3) a line containing v has at least $T(d-1)/d$ vertices initially open. In this case v may become open by time 2.

In this section we show that if p is sufficiently small then with high probability all of the vertices that are eventually open satisfy a condition like one of the three above. By doing this we prove a lower bound on the threshold function for a plane to become open.

Let Plane Active be the event that some plane becomes open. Let A be some integer, $1 \leq A \leq T$, which we will specify later. Let E be the event that there exists a vertex v such that:

(1) v is initially not open,
(2) the neighborhood of v has at most A vertices initially open, and
(3) each line containing v has at most $A/2$ vertices initially open.
(4) v becomes open.

Our goal is to show that for a value of p (to be specified later) that $\mathbb{P}(\text{Plane Active})$ is small. We do this by showing that both $\mathbb{P}(E)$ and $\mathbb{P}(\text{Plane Active} \setminus E)$ are small.

For each vertex v, let E_v be the event that v satisfies (1)-(4), and of such vertices, v is among the first to become open. If the event E occurs then one such vertex must become open at least as soon as all others, so $E \subseteq \cup_v E_v$, and by the union bound we have $\mathbb{P}(E) \leq n^d \mathbb{P}(E_v)$.

Lemma 6.1. *Let* $p = f(n)n^{-1-\beta}$ *with* $f(n) \to 0$ *as* $n \to \infty$, *and* $\beta = (\frac{2d^2}{T-A} + 1)\frac{2}{A}$. *Then* $\mathbb{P}(E) \to 0$ *as* $n \to \infty$.

Proof. We observed above that $\mathbb{P}(E) \leq n^d \mathbb{P}(E_v)$. Now if E_v occurs, by (2) at least $T - A$ vertices in the neighborhood of v must be initially closed but become open strictly before v; therefore, they violate at least one of (1)-(4). But since they start closed and become open, they must violate one of (2) or (3). By the pigeonhole principle, of the d lines through v, at least one must either contain $\frac{T-A}{2d}$ vertices which violate (2), or $\frac{T-A}{2d}$ vertices which violate (3).

The next two lemmas will show that these each happen with probability that is

$$o\left(n^{\frac{T-A}{2d}(1-\beta\frac{A}{2})}\right).$$

Rearranging the inequality $\beta > (\frac{2d^2}{T-A} + 1)\frac{2}{A}$ we see that $\mathbb{P}(E_v) = o(n^d)$ and

$$\mathbb{P}(E) \leq n^d \mathbb{P}(E_v) \to 0.$$

□

Lemma 6.2. *Let* $p = f(n)n^{-1-\beta}$ *with* $\beta \geq (\frac{2d^2}{T-A} + 1)\frac{2}{A}$. *Fix a line* L. *The probability that* L *has at least* $\frac{T-A}{2d}$ *vertices violating (2) is*

$$o\left(n^{\frac{T-A}{2d}(1-\beta\frac{A}{2})}\right).$$

Proof. If a given vertex violates (2), then its neighborhood excepting L must contain at least $A/2$ open vertices. These deleted neighborhoods are disjoint and the number of vertices that they have initially open are i.i.d. Bernoulli$((d-1)(n-1),p)$ random variables. The chance that a given one is at least $A/2$ is bounded by $c(np)^{A/2} = o\left(n^{-\beta A/2}\right)$. Then, the probability that at least $\frac{T-A}{2d}$ succeed is $o\left((n^{1-\beta A/2})^{\frac{T-A}{2d}}\right)$. $\square$

Lemma 6.3. *Let $p = f(n)n^{-1-\beta}$ with $\beta \geq (\frac{2d^2}{T-A}+1)\frac{2}{A}$. Fix a line L through v and suppose L has at most $A/2$ vertices initially open. The probability that L has at least $\frac{T-A}{2d}$ vertices violating (3) is*

$$o\left(n^{\frac{T-A}{2d}(1-\beta\frac{A}{2})}\right).$$

Proof. The probability that a given line has at least $A/2$ points initially open is bounded above by

$$\binom{n}{A/2}p^{A/2} \leq cf(n)^{A/2}n^{-\beta A/2} = o\left(n^{-\beta A/2}\right).$$

So of the $n(d-1)$ trials, the chance that at least $\frac{T-A}{2d}$ succeed $o\left((n^{1-\beta A/2})^{\frac{T-A}{2d}}\right)$. $\square$

Lemma 6.4. *Let $p = n^{-1-\beta}$. If $\beta > 0$ then $\mathbb{P}($Plane Active $\setminus E) \to 0$ as $n \to \infty$.*

Proof. There are $\binom{d}{2}n^{d-2}$ planes and Plane Active $= \cup_P(P$ becomes open$)$, so we have

$$\mathbb{P}(\text{Plane Active} \setminus E) \leq \binom{d}{2}n^{d-2}\mathbb{P}(\text{Plane Active}_P \setminus E).$$

Now if P becomes open but E does not occur, then since each point in P becomes open, they must all violate one of (1),(2), or (3). By the pigeonhole principle, at least $n^2/3$ of these points must together violate a single condition. We will check that the probability of these three cases each converge to 0 (even when multiplied by the n^{d-2} mentioned above):

- $n^2/3$ vertices are initially open

  The probability of this is bounded by

  $$\binom{n^2}{n^2/3}p^{n^2/3} \leq 2^{n^2}(p^{1/3})^{n^2} = o\left(n^{d-2}\right).$$

- $n^2/3$ vertices each are in a line with $A/2$ points initially open,

  This implies that there are at least $n/6$ parallel lines with at least $A/2$ points initially open. The probability of this is at most

  $$2\binom{n}{n/6}\left(\binom{n}{A/2}p^{A/2}\right)^{n/6} \leq 2\cdot 2^n\left(n^{-\beta A/2}\right)^{n/6} \leq 2\cdot 2^n n^{-n/6} = o\left(n^{d-2}\right).$$

- $n^2/3$ vertices each have a neighborhood with A vertices initially open,

  If a vertex v' has at least A vertices in its neighborhood initially open then either one of the two lines through v' in P contain at least $A/4$ vertices initially open or the crosslines through v' (not in P) contain at least $A/2$. This implies that there are at least $n/12$ parallel lines with at least $A/4$ vertices initially open or at least $n^2/6$ vertices with at least $A/2$ vertices in the cross lines out of P. The probability of the first is at most

  $$2\binom{n}{n/12}\left(\binom{n}{A/4}p^{A/4}\right)^{n/12} \leq 2\cdot 2^n\left(n^{-\beta A/4}\right)^{n/12} \leq n^{-cn} = o\left(n^{d-2}\right)$$

  and the second has probability at most

  $$\binom{n^2}{n^2/6}\left(\binom{dn}{A/2}p^{A/2}\right)^{n^2/6} \leq c\cdot 2^{n^2}\left(n^{-\beta A/2}\right)^{n^2/6} \leq n^{-cn^2} = o\left(n^{d-2}\right).$$

Thus $\mathbb{P}($Plane Active $\setminus E) \to 0$. $\square$

Proof of Theorem 2.3. To get the lower bound set $A = \lfloor T - \sqrt{T} \rfloor$ in Lemma 6.1. The lemma applies for

$$\beta = \frac{2}{T} + \frac{4d^2+2}{T^{3/2}}.$$

To get the upper bound set $\alpha = \left\lfloor \frac{\sqrt{T}}{2} \right\rfloor$ in Theorem 5.7. Then we get for

$$\beta < \frac{2}{T} + \frac{1}{T^{3/2}}$$

the probability of finding a plane eventually open is approaching 1. □

References

[1] Joan Adler and Uri Levi. Bootstrap percolation: visualizations and applications. *Brazilian Journal of Physics*, 33(3), 2003.

[2] M. Aizenman and J. L. Lebowitz. Metastability effects in bootstrap percolation. *J. Phys. A*, 21(19):3801–3813, 1988.

[3] József Balogh, Béla Bollobás, and Robert Morris. Bootstrap percolation in three dimensions. *Ann. Probab.*, 37(4):1329–1380, 2009.

[4] József Balogh, Hugo Bollobás, Béla amd Duminil-Copin, and Robert Morris. The sharp threshold for bootstrap percolation in all dimensions. *arXiv:1010.3326v2.*

[5] R. Cerf and F. Manzo. The threshold regime of finite volume bootstrap percolation. *Stochastic Process. Appl.*, 101(1):69–82, 2002.

[6] Raphaël Cerf and Emilio N. M. Cirillo. Finite size scaling in three-dimensional bootstrap percolation. *Ann. Probab.*, 27(4):1837–1850, 1999.

[7] J Chalupa, P L Leath, and G R Reich. Bootstrap percolation on a Bethe lattice. *J. Phys. C*, 12(1):L31–L35, 1979.

[8] Alexander E. Holroyd. Sharp metastability threshold for two-dimensional bootstrap percolation. *Probab. Theory Related Fields*, 125(2):195–224, 2003.

[9] Alexander E. Holroyd, Thomas M. Liggett, and Dan Romik. Integrals, partitions, and cellular automata. *Trans. Amer. Math. Soc.*, 356(8):3349–3368 (electronic), 2004.

[10] Aernout C. D. van Enter. Proof of Straley's argument for bootstrap percolation. *J. Statist. Phys.*, 48(3-4):943–945, 1987.